

## Study on Mouth Gymnastics for Orbicularis Oris Muscle

Akira Suzuki<sup>1,a</sup>, Nobuaki Nakazawa<sup>1,b</sup>, Yuuki Tanaka<sup>1,c</sup>,  
Yusaku Fujii<sup>2,d</sup>, Takao Yamaguchi<sup>2,e</sup>, and Toshikazu Matsui<sup>1,f</sup>

<sup>1</sup>29-1 Hon-Cho, Oota, Gunma, 373-0057 Japan

<sup>2</sup>1-5-1 Tenjin-Cho, Kiryu, Gunma, 376-8515 Japan

<sup>a</sup> <t12802202@gunma-u.ac.jp>, <sup>b</sup> <n.nakazawa@gunma-u.ac.jp>, <sup>c</sup> <ytanaka@gunma-u.ac.jp>,  
<sup>d</sup> <fujii@gunma-u.ac.jp>, <sup>e</sup> <yamagme3@gunma-u.ac.jp>, <sup>f</sup> <matsui@gunma-u.ac.jp>

**Keywords:** Mouth, Zygomaticus major and minor muscles, Orbicularis oris muscle, EMG, APDF

**Abstract.** There is a tendency that the elderly persons decline in feeding and swallowing. It is known that the oral training such as a facial exercise has an effect to facilitate a saliva secretion which is useful for the smoothness swallowing motion. This paper investigated the effective mouth gymnastics in the oral training practices, based on the biological signal of the electromyography (EMG). Here, we focused on the zygomaticus major and minor muscles, and orbicularis oris muscle for an evaluation. The obtained datum of EMG were analyzed by the amplitude probability distribution function (APDF) method to evaluate the effective gymnastics. As a result, it could be confirmed that the motion of puffing the cheeks and the vertical movement of the mouth corner had an effect to enhance the activity of the mouth muscles.


### 1. Introduction

Motion to swallow food and a drink are called "deglutition". The decline of oral sensation and the salivation with aging tends to decrease the function of the deglutition. As for some elderly, food and a drink are clogged up to the bronchus and may cause pneumonia. In the serious case, the nourishment is taken through a pipe, and the pleasure of the original meal will be lost by a nasal cavity. In order to facilitate a smoothness swallowing, it is known that the dysphagia by aging can be prevented by training oral neighborhood source and face exercises [1,2]. In addition, the mouthpiece-formed product to train oral neighboring sources has been developed [3]. The author has investigated the facial feature points such as the lips in the oral training practices for the purpose of the prevention of dysphagia [4]. About the effect of face exercises, the inspection is formed so far by quantity of salivation and opening degree, repetition saliva deglutition test [1], lips closedown power and syllable alternation repetition campaigns [2]. However, little work has been done on the oral practices based on the biological signal. Here, we used the biological signal of the electromyography (EMG) to evaluate the effective mouth gymnastics. The zygomaticus major and minor muscles, and orbicularis oris muscle were focused to consider the optimal practice, and the effective gymnastic was clarified by the amplitude probability distribution function (APDF) analysis.

### 2. Muscular activity of face practices

#### 2.1 Measurement

With reference to the exercises for the purpose of the improvement of the oral cavity function, namely, mobility and activity, there are the exercise of neck and shoulder, opening and shutting of the mouth, lips and tongue, cheeks blow up, and utterance. Here, we paid attention to the movement



a. Zygomaticus minor muscle  
 b. Zygomaticus major muscle  
 c. Orbicularis oris muscle

Fig.1 Measurement points of EMG




Fig.2 Scene of the oral training




Fig.3 Time series of data of EMG

of lips and cheeks from face exercise and measured the EMG signal. As shown in Fig.1, the electrodes were put near orbicularis oris muscle and zygomaticus major muscle and zygomaticus minor muscle, a ground was attached to a wrist. The electrode unit (Biometrics SX230) of the dry type was used for a measurement. The obtained EMG signal was taken to the personal computer with sampling frequency of 600[Hz].

## 2.2 Face practices

Fig.2 shows the lip and cheek postures when the face exercises were carried out. This figure includes three states from normal state to each state, pouting with saying “Uh...”(Motion A), spreading the corner of the mouth with saying “Ye...”(Motion B), and inflating the cheek (Motion C). In the practice of Motion A and Motion B, the muscle around the mouth is strengthened and it is effective in preventing food from running it down from a mouth. As for the practice of Motion C, it is intended to strengthen the muscle of the circumference of the mouth and the throat.

## 2.3 EMG signal and APDF analysis

Typical time trajectory of Motion A, Motion B, and Motion C are shown in Fig.3. The EMG signal of the zygomaticus major and minor muscles and the orbicularis oris muscle are assigned to Ch1 and Ch2, respectively. This evaluation is very often used [5,6]. In the exercise to pout over (Motion A), shown in Fig.3 (a), the orbicularis oris muscle is active while there are few changes in the activity of the zygomaticus major and minor muscles. In the exercise to widen the corners of the mouth (Motion B), shown in Fig.3 (b), both muscle parts are active. With reference to the exercise to inflate the cheek (Motion C), shown in Fig.3 (c), the amplitude level of the orbicularis oris muscle is larger, compared with Motion A and B. Here, the amplitude probability distribution function (APDF) of the obtained data was computed. It is used for an evaluation of a muscle burden [5,6]. Fig.4 shows




Fig.4 APDF analysis






Fig.5 APDF example of three practices



(a) Open width

(b) Horizontal movement of corner



(c) Vertical movement of corner

Fig.6 APDF of the mouth excises

the enlarged view of the time trajectory of EMG signal. Let  $T_i$  be the  $i$  th appearance time duration when the amplitude of the EMG signal became more than  $x\%$  of the EMG maximal voluntary contraction (MVC), the value of APDF can be expressed by the following equation.

$$P = \frac{1}{T_{total}} \sum_{i=1}^N T_i \times 100 \quad (1)$$

Were,  $N$  is the total number of the time duration  $T_i$  during the mouth gymnastics, and  $T_{total}$  is the total time duration and. Here, we derived the maximum amplitude of the EMG signal from whole trials in individual, which was set to the MVC. Fig.5 shows the APDF example of three practices. In this figure, the typical data of the orbicularis oris muscle was picked up. Note that the activity of the muscle can be regarded as higher if the percentage with respect to  $P=0.5$  is higher. From this figure, it is found that the percentage of Motion C with respect to  $P=0.5$  is higher than other motions and the exercise of inflating the cheek is effective for the training of the orbicularis oris muscle.

## 2.4 Mouth exercises

Next, we investigated the knack of the mouth excises for effective training. Fig.6 shows the experimental results of changes in the mouth open width, horizontal and vertical movement of the mouth corner. The excise of opening mouth (Fig.6(a)) is relatively small, compared with the excise of moving the mouth corner (Fig.6(b) and (c)). In this case, the zygomaticus muscle is regarded as low active. In Fig.6(b) and (c), it can be confirmed that the activity of both the zygomaticus muscle and the orbicularis oris muscle are active and it becomes a higher activity as the movement of the mouth corner is larger. Especially, the vertical movement of the mouth corner has a higher percentage with respect to  $P=0.5$  and it can be considered as an effective for the oral training.

## 3 Conclusions

In this study, we investigated the face exercises through the measurement of the EMG around the oral cavity. Here, the zygomaticus major and minor muscles, and orbicularis oris muscle were focused for an evaluation by using APDF analysis. As a result, it could be found that the motion of puffing the cheeks and the vertical movement of the mouth corner had an effect to enhance the muscle activity of the mouth. As a future works, we would like to construct the mouth training GUI based on the effect practices.

## Acknowledgements

This research was partially supported by the Ministry of Education, Science, Sports and Culture, Grant-in-Aid for Scientific Research (C), 2015-2017, No.15K05888.

## References

- [1] T.Ooka, T. Haino, S. Hironaka and Y. Mukai, The Effect of Daily Oral Function Training in the Elderly, *Journal of dental health*, Vol.58, No.2, pp.88-94, 2008.
- [2] T.Kikuchi, T.Yoneyama, T.Teshima, F.Hori uchi, K.Miyatake, M.Adachi, M.Ishida and K.Nishiwaki, Effect of Oral Function Training and Feeding Assistance Care for Nutritional Improvement in the Elderly, *Japanese journal of gerodontology*, Vol.20, No.3, 208-213, 2005.
- [3] <http://www.combiwellness.co.jp/cata/yobo/labilin.html>J.
- [4] A. Suzuki, N.Nakazawa, Y.Tanaka, Y.Fujii, T.Yamaguchi, and T.Matsui, Effective Mouth Gymnastics for Orbicularis Oris Muscle, *Proc. ICTSS 2017*, May 2017.
- [5] T. Mukai, H.Oyama, K. Noro, The evaluation of muscle activities during use a ballpoint pen by using APDF analysis, *Japan Society of Physical Anthropology*, Vol.11, No.3, pp. 13-19, 2006.
- [6] K. Tomioka, Study on Myoelectric APDF for Consumer Product Development and Evaluation (In Japanese), *Toshiba Review*, Vol.56, No.6, pp. 58-61, 2001.