

## Early Effects of Streptozotocin Diabetes on Capillaries in Spinal Cord

Kanokporn Plaengrit<sup>1, a</sup>, Sirinush Sricharoenvej<sup>1, b</sup>, Passara Lanlua<sup>1, c</sup>,  
Apichaya Niyomchan<sup>1, d</sup>, and Amornrat Chookliang<sup>1, e</sup>

<sup>1</sup> Department of Anatomy, Faculty of Medicine Siriraj Hospital, Mahidol University, Bangkoknoi, Bangkok10700, Thailand

<sup>a</sup><kanokkhaw@gmail.com>, <sup>b</sup>< sirinush.sri@mahidol.ac.th>, <sup>c</sup><passara.lan@mahidol.ac.th>

<sup>d</sup><apichaya.niy@mahidol.ac.th>, <sup>e</sup><tongzaanatomy@gmail.com>

**Keywords:** diabetic microangiopathy, neuronal complications, spinal capillary

**Abstract.** Diabetic microangiopathy is a well-known cause of neuronal complications mainly in peripheral nerves and brain. Additionally, it also strongly associates with the development of spinal infarction followed by the physical disability to seriously affected the quality of life in diabetic patients. However, only few study is known about the alterations of the spinal capillary in diabetes. Therefore, this study aimed to illustrate morphological changes of spinal capillaries in the early stage of streptozotocin (STZ)- induced diabetic rat compared with control. In diabetes, the endothelial cell and pericyte became hypertrophy. Under the ultrastructural level, the endothelial cell contained a vast number of ribosomes and vesicles, enlarged mitochondria, and disrupted tight junction, while the vascular pericyte had numerous free ribosomes and a pair of centrioles. Furthermore, basement membrane thickening was detected. This basic knowledge will be beneficial for raised awareness and receiving in early diagnosis of the spinal vascular disorder in diabetic patients.

### 1. Introduction

During diabetes mellitus (DM), a very sensitive target to high glucose is the blood vessel. Hyperglycemia initiates the pathogenesis of macroangiopathy and microangiopathy [1], causing insufficient blood-induced dysfunctions of the several organ systems. Many studies have investigated the alterations of capillary components; endothelial cell and pericyte in the peripheral tissues and the nervous tissues. In the general studies of diabetic complications, STZ has been used for producing a suitable hyperglycemic animal model. The STZ is highly preferential toxicity to the pancreatic beta cells that subsequently undergo cell necrosis causing diabetes. Moreover, the pathophysiological characterizations of this model can be compared with those in human with diabetes [2, 3]. In the early stage of diabetes, the retinal capillary cell death caused retinopathy [4], as well as the glomerular endothelial cell proliferation with thickened basement membrane occurred in relation to progressing histological lesion in nephropathy [5]. Additionally, the sinusoidal capillaries observing in hepatic microvascular casts are decreased in diameter [6]. Regarding the severe diabetic neuropathy, the nerve capillary pathology has also been studied, for instance, a decrease in endoneurial capillary density, an increase in thickening of total diffusion barrier and a reduction of oxygen diffusing capacity [7]. In the cerebral microvessels, the vascular lumen stenosis, thickened vascular basement membranes, endothelial cell swelling, and the shedding has been revealed in the STZ- induced diabetic rats [8]. Furthermore, one cause of disabling event in the patient with diabetes have also been reported with the abnormal vascular wall and the embolic occlusion in a vessel leading to an inadequate blood circulation and subsequent degenerative changes in the spinal cord. As the spinal cord is the center route to carry nerve impulses, the clinical presentations commonly

relate to the sensory deficit, progressive weakness symptoms, and autonomic control disturbance, corresponding to the levels of the cord lesion [9, 10]. However, few studies examine spinal capillaries, especially the architecture of diabetic vasculopathy. Then, this study focused on the histological and ultramicroscopic examinations of spinal capillaries in the short duration STZ-induced DM.

## **2. Materials and Methods**

### **2.1 Animal preparation**

Fifteen male Sprague-Dawley rats (5-8 week old, 200-270 g body weight) were provided by the National Laboratory Animal Center, Mahidol University, Salaya, Nakornpathom, Thailand. Rats were taken care in accordance with the Mahidol University Care and Use of Laboratory Animal.

### **2.2 Experiment protocols**

Rats were randomly divided into two groups: control (n=6) and diabetic animals (n=9). Type 1 diabetes was induced by a single intraperitoneal injection of STZ (60 mg/kg body weight) dissolved in citrate buffer (pH 4.5). The control rats received only the citrate buffer. After STZ administration for 48 and 72 hours, the blood samples were collected from the tail vein and the blood glucose level was measured by a glucose meter (OneTouch® Ultra®, Colorado, USA). Rats with  $\geq 300$  mg/dL blood glucose level with symptoms of polyuria, polyphagia, and polydipsia were considered as diabetes and included in this study. At the end of the experiment (4 weeks after the induction), the animals were measured the blood glucose levels before sacrifice and anesthetized by halothane inhalation. Whole animal perfusion through the heart was processed for histological and electron microscopic examinations.

### **2.3 Histological study**

The 0.9% NaCl perfusion was performed, followed by injection of Bouin's solution to preserve tissues. Next, the spinal cords, which were separated into cervical enlargements (C), thoracic segment (T), lumbosacral enlargements (L), were harvested and immersed in the same fixative overnight. After rinsing with 70% ethanol several times, these specimens were dehydrated in a graded series of ethanol, cleared in xylene, infiltrated, and embedded in paraffin. Serial cross sections were cut with a microtome (Leica RM 2035, Wetzlar, Germany) at 6  $\mu$ m thickness, mounted on glass slides and stained with the routine hematoxylin and eosin (H&E) technique. Photography and morphometric studies were done under a light microscope (LM) (Axiostar plus, Jena, Germany) connected to a digital camera (AxioCam MRc, Jena, Germany).

### **2.4 Transmission electron microscopy**

The 0.1 M phosphate buffered saline (PBS) solution was perfused through the heart and subsequently injected with 2.5% glutaraldehyde in 0.1 M PBS to fix the tissues. The spinal cords including C, T, and L were removed and cut into small cubes. The tissue was postfixed in 0.1% osmium tetroxide, dehydrated in ethanol series, infiltrated in propylene oxide, and embedded in the plastic. The ultrathin sections were cut with an ultramicrotome (Leica EM UC6, Vienna, Austria), mounted on copper grids, counterstained with 1% acetate and lead citrate, and observed under the JEOL JEM 100S transmission electron microscope equipped with a digital camera (Tokyo, Japan).

### 3. Results

#### 3.1 Histology of spinal capillary

Similar alterations during early STZ diabetic stage were showed in the spinal capillaries in each area of gray and white matters from all spinal levels (C, T, and L). Under the LM, the spinal cords consisted of neurons, nerve fibers, neuroglia, and capillaries. Concerning to the spinal capillaries, they occupied among neurons and neuroglia in the gray matter, while they paralleled to nerve fibers and neuroglia in the white matter. Each spinal capillary was formed by two cell types; endothelial cell and pericyte. The endothelial cell had a flat basophilic nucleus enclosed spherical-shaped capillary lumen. The prominent nucleus of pericytes capped on the basal side of endothelial cell layer (Figs. 1A, 1C). In the STZ-induced diabetic rats, both endothelial cells and pericytes contained larger nuclei than those in the control rats. (Figs. 1B, 1D).

#### 3.2 Ultrastructure of spinal capillary

In the ultramicroscopic analysis of normal capillaries, the endothelial cell had a single flattened endothelial nucleus and a thin layer of cytoplasm surrounded a large round capillary lumen. Endothelial nucleus contained electron-lucent euchromatin and electron-dense heterochromatin (Fig. 2A). A large number of mitochondria inside the spinal endothelial cell were visible as a rod or spherical shape bounded with the double membrane layers. Rough endoplasmic reticulum (rER) displayed the slimline cisternae attached by numerous dark spots of ribosomes. Between endothelial cell and pericyte, the basal surface of the endothelial cell was underlined by the thin fibrillar layer of basement membrane (Fig. 2B). In the STZ diabetic capillaries, there were numerous free ribosomes scattered within a large volume of cytoplasm together with a group of short rER with dark stained elements. Basement membrane thickening in capillaries was demonstrated in figure 3A. A marked enlargement of mitochondria with cristae disruption was observed (Fig. 3B). Moreover, there were a large vacuole and a lot of vesicles in the cytoplasm near the basement membrane and subplasmalemmal zone (Figs. 3A-C). Apart from the endothelial cell, pericyte was also defined as the vascular cells. It became enlarged in diabetes and contained a large number of electron-dense spots of free ribosomes, particularly in the perinuclear region. The long cisternae of rER and lightly stained mitochondrial matrix with a few cristae also presented. In addition, a pair of tube-shaped centrioles appeared near the nuclear membrane (Fig. 3D).




Fig. 1. Light micrographs of the spinal capillaries in gray (1A, 1B) and white matters (1C, 1D) of the spinal cords in control (1A, 1C) and diabetic (1B, 1D) rats. Code: E= endothelial cell, P= pericyte, N= neuron, O= oligodendrocyte, A= astrocyte. H&E staining.

In the endothelial tight junction (TJ), the electron-dense line on the plasma membrane of adjacent cells was structurally intact in the normal condition (Fig. 4A). During diabetes, the alteration of TJ was the imperfect anchoring of the unit membrane, leading to the small gap formation between two adjacent endothelial cells. Additionally, a reduced electron density of the TJ site occurred in early STZ-induced diabetes (Fig. 4B).




Fig. 2. Transmission electron micrographs of the normal capillary at low (2A) and high (2B) magnifications. Code: L= capillary lumen, Nu= endothelial nucleus, Ax= axon, E= endothelial cell, m= mitochondria, r= ribosomes, rER= rough endoplasmic reticulum, B= basement membrane, P= pericyte.




Fig. 3. Cellular injuries of the STZ diabetic spinal capillaries were found in endothelial cells (3A-C) and pericyte (3D). Code: E= endothelial cell, rbc= red blood cell, r= ribosomes, a dashed arrow= dense material group of short rough endoplasmic reticulum, Ve= a lot of endocytic vesicles, a two-headed arrow= thickened basement membrane Nu= endothelial nucleus, B= basement membrane, white stars= dilated mitochondria, Va= a large vacuole, L= capillary lumen, P= pericyte, m= mitochondria, rER= rough endoplasmic reticulum, mi= microfilaments, Cen= a pair of centrioles.




Fig. 4. Transmission electron micrographs of tight junction interlocking between endothelial cells of the spinal capillary in the control (4A) and STZ- induced diabetic (4B) rats.

Code: E= endothelial cell, Nu= endothelial nucleus,  
 TJ= intact electron dense line of tight junction, B= basement membrane,  
 L= capillary lumen, a black arrowhead= detachment of TJ.

#### 4. Discussion

The development of diabetic microvascular complications in the spinal cord occurred in the early phase of STZ diabetes. The endothelial cell lining of the capillary lumen and the pericyte wrapping around the endothelial cell of the capillaries increased in their sizes. The excess glucose in these cells during diabetes can be converted into the sorbitol and fructose via the polyol pathway contributing to an increase in intracellular oncotic pressure [11]. This change results in water entering the cell causing cellular edema. Moreover, the high glucose level can induce transcellular transportation. During diabetes, advanced glycation end product (AGE) induces an increase in vascular endothelial growth factor (VEGF) expression through nuclear factor kappa B (NF- $\kappa$ B) pathway. Then, VEGF can increase the endothelial permeability via an endothelial nitric oxide synthase (eNOS)- the dependent process of transcytotic transport in plasma membrane caveolae through a mechanism that involves the phosphoinositide 3 kinase (PI3K)- protein kinase B (Akt)- eNOS signaling [12]. Therefore, the formation of numerous interconnected membrane invaginations, termed the vesicles and large vacuoles, was also found in the cytoplasm of diabetic endothelial cells. Furthermore, an increased VEGF robustly activates the PI3K-Akt pathway. The phosphorylation of the Akt targets contributes to cell survival, growth, and proliferation. Thus, abundant free ribosomes and duplicated short rER, that were essential for protein synthesis during the interphase of the cell cycle, were observed in the cytoplasm of the diabetic endothelial cell. In addition, a pair of centrioles for cell division was seen in the pericyte near its nucleus. It was concluded that AGE in hyperglycemic environment stimulates aberrant angiogenesis through the angiogenic peptide VEGF [13]. VEGF directly contributes to capillary basement membrane thickening via increased type IV collagen deposition in diabetes [14]. Moreover, the glycation products themselves can induce gene expression and synthesis of tumor necrotic factor alpha (TNF- $\alpha$ ) in the endothelial cells [15]. The TNF- $\alpha$  represses the claudin 5, an integral membrane protein which is a critical component of endothelial TJ, via protein kinase C/NF- $\kappa$ B signaling pathway in diabetic condition [16]. Thereby, the detached TJ occurred in the STZ diabetic rat, implying a vascular permeability defect. The reported alterations of spinal capillaries in early stage of STZ-induced diabetes in this study should persist to fully elucidate the short-term effects of diabetes type 1 on spinal microvascular dysfunction in order to better early therapeutic detection to reduce chronic diabetic complications.

## 5. Conclusion

Early developed STZ diabetes causes the spinal capillary changes including the endothelial cell lining of the capillary lumen and the pericyte wrapping around the endothelial cell. The capillary abnormalities may be suggested as a possible pathogenic cause of diabetic neuropathy in latter. An early detection of spinal capillary lesion gives the chance to reduce the severity of microangiopathy related to neuropathy in the diabetic patients.

## Acknowledgements

This study was supported by the Siriraj Graduate Scholarships, Siriraj Research Fund, and Chalermphrakiat Grant, Faculty of Medicine Siriraj Hospital, Mahidol University, Bangkok, Thailand.

## References

- [1] WT. Cade, "Diabetes-related microvascular and macrovascular diseases in the physical therapy setting", *Phys. Ther.*, Vol. 88, No. 11, pp. 1322-1355, 2008.
- [2] P. Lanlua, Y. Tongpob, A. Niyomchan, I. Phoungpetchara, and S. Sricharoenvej, "Impairments of renal morphology in short-termed streptozotocin-induced diabetic rat", *Siriraj Med. J.*, Vol. 59, suppl. 2, pp. S149-S151, 2007.
- [3] S. Sricharoenvej, Y. Tongpob, P. Lanlua, S. piyawinijwong, J. Roongruangchai, and I. Phoungpetchara, *J. Med. Assoc. Thai.*, Vol. 9 No. 12, pp. 2677-2682, 2007.
- [4] RA. Kowluru and S Odenbach, "Effect of long-term administration of  $\alpha$ -lipoic acid on retinal capillary cell death and the development of retinopathy in diabetic rats", *Diabetes*, Vol. 53, No. 12, pp. 3233-3238, 2004.
- [5] T. Nakagawa, W. Sato, O. Glushakova, M. Heinig, T. Clarke, M. Campbell-Thompson, Y. Yuzawa, MA. Atkinson, RJ. Johnson, and B. Croker, "Diabetic endothelial nitric oxide synthase knockout mice develop advanced diabetic nephropathy", *J. AM. Nephrol.*, Vol. 18, No. 2, pp. 539-550, 2007.
- [6] S. Baimai, A. Niyomchan, P. Lanlua, J. Roongruangchai, and S. Sricharoenvej, "Modification of hepatic microvasculature in chronic diabetic rat", *Siriraj Med. J.*, Vol. 61, suppl. 1, pp. S5-S8, 2009.
- [7] RA. Mailik, PG. Newrick, AK. Sharma, A. Jennings, AK. Ah-See, TM. Mayhew, J. Jakubowski, AJ. Boulton, and JD. Ward, "Microangiopathy in human diabetic neuropathy: relationship between capillary abnormalities and the severity of neuropathy", *Diabetologia*, Vol. 32, No. 2, pp. 92-102, 1989.
- [8] H. Yang, S. Fan, D. Song, Z. Wang, S. Ma, S. Li, X. Li, M. Xu, M. Xu, and X. Wang, "Long-termed streptozotocin-induced diabetes in rats leads to severe damage of brain blood vessels and neurons via enhanced oxidative stress", *Mol. Med. Rep.*, Vol. 7, No. 2, pp. 431-440, 2013.
- [9] F. Romi and H. Naess, "Characteristics of spinal cord stroke in clinical neurology", *Eur. Neurol.*, Vol. 66, No. 5, pp. 305-309, 2011.
- [10] T. Sugihara, K. Kido, Y. Sasamori, M. Shiba, and T. Ayabe, "Spinal cord infarction in diabetic pregnancy: a case report", *J. Obstet. Gynaecol. Res.*, Vol. 39, No. 10, pp. 1471-1475, 2013.
- [11] JR. Lemasters, T. Qian, CA. Bradham, DA. Brenner, WE. Cascio, LC. Trost, Y. Nishimura, AL. Nieminen, and B. Hermam, "Mitochondrial dysfunction in the pathogenesis of necrotic and apoptotic cell death", *J. Bioenerg. Biomembr.*, Vol. 31, No. 4, pp. 305-319, 1999.
- [12] J. Chen, F. Braet, S. Brodsky, T. Weinstein, V. Romanov, E. Noiri, and MS. Goligorsky, "VEGF-induced mobilization of caveolae and increase in permeability of endothelial cells", *Am. J. Physiol. Cell Physiol.*, Vol. 282, No. 5, pp. 1053-1063, 2002.

- [13] R. Humar, FN. Kiefer, H. Berns, TJ. Resink, and EJ. Battegay, "Hypoxia enhances vascular cell proliferation and angiogenesis in vitro via rapamycin (mTOR)-dependent signaling", *FASEB J.*, Vol. 16, No. 8, pp. 771-780, 2002.
- [14] EJ. Kuiper, JM. Hughes, RJ. Van Geest, IM. Vogels, R. Goldschmeding, CJ. Van Noorden, RO. Schlingemann, and I. Klaassen, "Effect of VEGF A on expression of profibrotic growth factor and extracellular matrix genes in the retina", *Invest. Ophthalmol. Vis. Sci.*, Vol. 48, No. 9, pp. 4267-4276, 2007.
- [15] G. Rashid, S. Benchetrit, D. Fishman, and J. Bernheim, "Effect of advanced glycation end-products on gene expression and synthesis of TNF-alpha and endothelial nitric oxide synthase by endothelial cells", *Kidney Int.*, Vol. 66, No. 3, pp. 1099-1106, 2004.
- [16] CA. Aveleria, C. Lin, SF. Abcouwer, AF. Ambrosio, and DV. Antonetti, "TNF- $\alpha$  signals through PKC $\zeta$ /NF- $\kappa$ B to alter the tight junction complex and increase retinal endothelial cell permeability", *Diabetes*, Vol. 59, No. 11, pp. 2872-2882, 2010.